

SPECIFICATION

TITLE OF THE INVENTION

PROCESS FOR PREPARING GLYCIDYLPHTHALIMIDE

5

BACKGROUND OF THE INVENTION

The present invention relates to a process for preparing glycidylphthalimide, especially optically active glycidylphthalimide, which is useful as an intermediate for preparing medicines, agrochemicals or physiologically active compounds.

PRIOR ART

Glycidylphthalimide, especially optically active glycidylphthalimide is utilizable as an important intermediate for preparing many kinds of optically active medicines.

Generally, optically active medicines and intermediates thereof are required of 98% e.e. with optical purity. Therefore it is very important to establish the process for preparation of optically active glycidylphthalimide having optically high purity easily.

As general methods for preparing racemic glycidylphthalimide, a method for preparing it by refluxing potassium phthalimide in racemic epichlorohydrin solvent (J.

25

Am. Chem. Soc. Vol. 117, p 11220-11229(1995): abbreviated
as Reference 1) and a method for preparing it by reacting
an equal molar quantity of potassium phthalimide and a
racemic epihalohydrin in DMF (Helv. Chim. Acta Vol. 73, p
5 912-915(1990): abbreviated as Reference 2) have been
developed. Furthermore, a method for preparing it by
reacting phthalimide and racemic epichlorohydrin in tetra-
n-butylammonium iodide and potassium carbonate in a
microwave oven is described in Synlett p 873-874(1996)
10 (abbreviated as Reference 3).

On the other hand in regard to methods for preparing
optically active glycidylphthalimide, a method for
preparing it by following steps (a) to (c): (a) refluxing
phthalimide in optically active epichlorohydrin solvent to
15 obtain optically active N-(3-chloro-2-
hydroxypropyl)phthalimide, (b) isolating and purifying the
compound and then (c) cyclizing the compound with sodium
hydride is described in US Patent No. 5608110. In WO
99/24393, (S)-glycidylphthalimide is described, but there
20 is no description on the method for preparing it.

According to the above methods for preparing racemic
glycidylphthalimide described in Reference 1, when
potassium phthalimide and an optically active epihalohydrin
in the amount of the solvent is reacted under refluxing,
25 optical purity of the epihalohydrin decreases and therefore,

optical purity of glycidylphthalimide produced therein decreases. When optically active epichlorohydrin (99%e.e.) and potassium phthalimide are reacted in DMF, racemization proceeds to give optically active glycidylphthalimide with
5 poor optical purity (63% e.e.) (See Coperative example 1).


On the other hand, the object compound is not obtained in good yield when the reaction is carried out in a microwave oven according to the method described in Reference 3. According to the method of US Patent No.
10 5608110, an intermediate, N-(3-chloro-2-hydroxypropyl)phthalimide must be isolated and the yield thereof and the object compound are not satisfactory.

The present inventors have extensively studied to solve the above problem and it have been found that by
15 reacting an optically active epihalohydrin and an alkali metal phthalimide in an alcohol as a solvent, optically active glycidylphthalimide is easily obtainable in good yield and with optically high purity. Furthermore, it has been found that by reacting a racemic epihalohydrin and an
20 alkali metal phthalimide, racemic glycidylphthalimide is also easily obtainable in good yield.


In addition, it has been found that racemic glycidylphthalimide and its optically active compound are easily obtainable in good yield and optically active
25 glycidylphthalimide is obtainable with optically high

purity, by reacting phthalimide and an epihalohydrin or an optically active epihalohydrin in an organic solvent in the presence of an alkali metal carbonate, an alkali metal hydrogencarbonate or a quaternary ammonium salt to obtain a
 5 N-(3-halogeno-2-hydroxypropyl)phthalimide and then by cyclizing the compound with an alkali metal alkoxide, too.


Namely the present invention relates to a process for preparing a glycidylphthalimide represented by the following formula (1):



which comprises reacting in an alcohol solvent an alkali metal phthalimide represented by the following formula (2):




wherein M is an alkali metal,
 15 with an epihalohydrin represented by the following formula (3):




wherein X is a halogen atom; or
 reacting phthalimide and an epihalohydrin (3) in an organic
 20 solvent in the presence of an alkali metal carbonate, an

alkali metal hydrogencarbonate or a quaternary ammonium salt (4) represented by the following formula:



wherein R_1 , R_2 , R_3 and R_4 are the same or different, C_{1-16} alkyl, C_{2-16} alkenyl, aryl-alkyl(C_{1-16}) or aryl, and X is chloro ion, bromo ion, iodo ion, hydrogensulfate ion or hydroxy ion,

to prepare a N-(3-halogeno-2-hydroxypropyl)phthalimide represented by the following formula (5):



wherein X is the same as defined above,

and then by cyclizing the compound (5) with an alkali metal alkoxide.

The present invention relates to a process for preparing glycidylphthalimide (1) which comprises reacting in an alcohol solvent an alkali metal phthalimide (2) with an epihalohydrin (This invention refers to process (A)).

Especially the present invention relates to a process for preparing optically active glycidylphthalimide (1) which comprises reacting an alkali metal phthalimide (2) with an optically active epihalohydrin (3) in an alcohol solvent.

The present invention also relates to the above mentioned process (A), wherein the process is carried out in the presence of a quaternary ammonium salt (4).

5 The present invention relates to a process for preparing glycidylphthalimide (1) which comprises reacting phthalimide and an epihalohydrin (3) in an organic solvent in the presence of an alkali metal carbonate, an alkali metal hydrogencarbonate or a quaternary ammonium salt (4) to prepare a N-(3-halogeno-2-hydroxypropyl)phthalimide (5)
10 and then by cyclizing the compound (5) with an alkali metal alkoxide (This invention refers to process (B)).

The present invention relates to a process for preparing glycidylphthalimide (1) which comprises carrying out the above process (B) without isolating a N-(3-halogeno-2-hydroxypropyl)phthalimide (5) prepared in the
15 first step, namely one pot method.

Especially the present invention relates to process (B) for preparing optically active glycidylphthalimide (1) which comprises reacting phthalimide with an optically
20 active epihalohydrin, especially optically active epichlorohydrin.

The present invention also relates to a process for preparing a N-(3-halogeno-2-hydroxypropyl)phthalimide (5) which comprises reacting phthalimide and an epihalohydrin
25 (3) in an organic solvent in the presence of an alkali

metal carbonate, an alkali metal hydrogencarbonate or a quaternary ammonium salt (4). Furthermore, the present invention relates to a process for preparing an optically active N-(3-halogeno-2-hydroxypropyl)phthalimide (5) which
5 comprises reacting phthalimide with an optically active epihalohydrin in an organic solvent.

The alkali metal phthalimide (2) used in the present invention relating to process (A) includes the compound in which an alkali metal is bound with nitrogen atom
10 consisting of an imido group of the compound, such as sodium phthalimide, potassium phthalimide, cesium phthalimide, and preferably potassium phthalimide.

The epihalohydrin (3) used in the present invention relating to process (A) includes epichlorohydrin, epibromohydrin and epiiodohydrin, preferably optically
15 active isomer thereof, especially preferably an optically active epihalohydrin. The amount of the epihalohydrin is 1 to 4 moles to the alkali metal phthalimide (2), preferably 2 to 3 moles.

20 The alcohol solvent used in the present invention relating to process (A) includes primary alcohols, such as methanol, ethanol, 1-propanol, 1-butanol, 1-pentanol, 1-hexanol, etc., secondary alcohols, such as isopropanol, 2-butanol, 2-pentanol, 3-pentanol, 2-hexanol, cyclohexanol,
25 2-heptanol, 3-heptanol, etc., tertiary alcohols, such as

tert-butanol, tert-pentanol, etc., preferably secondary or tertiary alcohols, more preferably isopropanol, 2-butanol and tert-butanol.

5 The amount of the alcohol solvent is preferably used 2 to 20 times (w/w) that of the alkali metal phthalimide (2).

The quaternary ammonium salt used in the present invention relating to process (A) is not limited, but includes benzyltrimethylammonium chloride, diallyldimethylammonium chloride, benzyltrimethylammonium bromide, n-octyltrimethylammonium bromide, stearyltrimethylammonium bromide, cetyltrimethylammonium bromide, tetra n-butylammonium iodide, β -methylcholinium iodide, tetra n-butylammonium hydrogensulfate, phenyltrimethylammonium hydroxide, etc.

15 The reaction relating to process (A) is promoted by addition of a quaternary ammonium salt (4) in the reaction system and therefore, the yield of glycidylphthalimide (1) or its optically active compound can be increased.

20 The amount of the quaternary ammonium salt (4) is catalytic amount to the alkali metal phthalimide (2), preferably 0.005 to 0.1 mole.

The reaction relating to process (A) is preferably carried out at -10 to 60°C , more preferably 0 to 30°C . When the reaction is carried out at below -10°C the reaction is restrained (controlled) and it is not desirable.

When the reaction is carried out beyond 60C°, the side reaction proceeds and therefore, it causes decrease of the yield on the object compound (1) and, in case that an epihalohydrin used is optically active, racemization of the compound proceeds and it causes decrease of optical purity of optically active glycidylphthalimide (1).

One of the merits of the present invention relating to process (A) is to that glycidylphthalimide (1), especially optically active glycidylphthalimide is obtained in good yield with highly optical purity by such a very simple procedure as removing the alcohol solvent and washing a solvent for extraction after the reaction is completed.

The solvent for extraction of the object compound is not limited as long as the compound is dissolved in said solvent, but includes acetates, such as methyl acetate, ethyl acetate, etc., and halogeno compounds, such as methylene chloride, chloroform, 1,2-dichloroethane, etc. The product may be, if necessary, purified by crystallization and column chromatography.

The epihalohydrin (3) used in the present invention relating to process (B) includes epichlorohydrin, epibromohydrin and epiodohydrin, preferably optically active form thereof, especially preferably an optically active epihalohydrin. The amount of the epihalohydrin is 1 to 3 moles to the alkali metal phthalimide (2), preferably

1 to 2 moles.

The alkali metal carbonate used in the present invention relating to process (B) is not limited, but includes lithium carbonate, sodium carbonate, potassium carbonate and cesium carbonate, and the alkali metal hydrogencarbonate used in the present invention relating to process (B) is not limited, but includes lithium hydrogencarbonate, sodium hydrogencarbonate, potassium hydrogencarbonate and cesium hydrogencarbonate. These carbonate salts may be hydrous or anhydrous, but in case preparing optically active glycidylphthalimide, an anhydrous carbonate salt is preferable. The amount of the alkali metal carbonate or the alkali metal hydrogencarbonate may be either stoichiometric amount or catalytic amount to phthalimide or, preferably 0.01 to 3 moles.

The quaternary ammonium salt used in the present invention relating to process (B) is the same as used in process (A).

The amount of the quaternary ammonium salt (4) is catalytic amount to the alkali metal phthalimide (2), preferably 0.005 to 0.1 mole. When the quaternary ammonium salt (4) is used together with the above alkali metal carbonate or alkali metal hydrogencarbonate, the reaction relating to process (B) is promoted to give the product in good yield.

The organic solvent used in the present invention relating to process (B) is not limited as long as it does not react with reactants (e.g., amines, carboxylic acid or a compound having glycidyl group), but in case preparing optically active glycidylphthalimide (1), alcohols, ethers and a mixture thereof are preferably used. The alcohol includes primary alcohols, such as methanol, ethanol, 1-propanol, 1-butanol, 1-pentanol, 1-hexanol, etc., secondary alcohols, such as isopropanol, 2-butanol, 2-pentanol, 3-pentanol, 2-hexanol, cyclohexanol, 2-heptanol, 3-heptanol, etc., and tertiary alcohols, such as tert-butanol, tert-pentanol, etc., preferably methanol, isopropanol and tert-butanol. The ether includes diethyl ether, dibutyl ether, tert-butyl methyl ether, 1,2-dimethoxyethane, 1,4-dioxane, tetrahydrofuran (THF), etc. The ratio of the mixture of the solvents is not limited. The amount of the solvent is preferably used 4 to 20 times (w/w) that of phthalimide.

The alkali metal alkoxide used in the present invention relating to process (B) (in 2nd step) includes sodium methoxide, potassium methoxide, sodium ethoxide, potassium ethoxide, sodium n-propoxide, potassium n-propoxide, sodium isopropoxide, potassium isopropoxide, sodium n-butoxide, potassium n-butoxide, sodium tert-butoxide, potassium tert-butoxide, sodium tert-amylate, potassium amylate, sodium n-hexylate, potassium n-hexylate,

etc.

The alkali metal hydroxide should be added after phthalimide and an epihalohydrin (3) are completely condensed to give a N-(3-halogeno-2-
5 hydroxypropyl)phthalimide. When the alkali metal alkoxide is added while phthalimide and an epihalohydrin is being condensed, the side reaction occurs and the yield of the object compound, especially the yield of the optically active compound decreases (See Coparative example 5). The
10 alkali metal hydroxide is preferably added in several divided portions or is preferably gradually added to the reaction mixture after resolving it in an alcohol or an ether.

The reaction relating to process (B) is preferably
15 carried out at -10 to 60°C , more preferably 0 to 50°C .

Furthermore, there are obtainable a N-(3-halogeno-2-hydroxypropyl)phthalimide (5) in good yield or its optically active compound in good yield with high optical purity by reacting phthalimide with an epihalohydrin (3) or
20 an optically active epihalohydrin (3) in the presence of an alkali metal carbonate, an alkali metal hydrogencarbonate or a quaternary ammonium salt (4) in an organic solvent, preferably an alcohol or an ether and then by extracting and purifying the product by crystallization or column
25 chromatography. Thus obtained N-(3-halogeno-2-

hydroxypropyl)phthalimide(5) is very useful for an intermediate of glycidylphthalate or other compounds.

One of the merits of the present invention relating to process (B) is to that the process is carried out by using
5 cheaper materials and cheaper solvents, and glycidylphthalimide (1), especially optically active glycidylphthalimide is obtained in good yield with highly optical purity by such a very simple procedure as the solvent is removed and the product is washed with the
10 solvent for extraction after the reaction is completed. The solvent for extraction of the object compound is not limited as long as the compound is dissolved in said solvent, but includes acetates such as methyl acetate, ethyl acetate, etc., and halogeno compounds, such as
15 methylene chloride, chloroform, 1,2-dichloroethane, etc. The product may be, if necessary, purified by crystallization and column chromatography.

Another of the merits of the present invention relating to process (B) is to that without isolation of a
20 N-(3-halogeno-2-hydroxypropyl)phthalimide(5) or its optically active compound obtained by reacting phthalimide with an epihalohydrin (3) or an optically active epihalohydrin, the product is reacted with an alkali metal alkoxide to give glycidylphthalimide (1) or its optically
25 activated compound. Using the one pot method the object

compound (1) and its optically active compound are obtained in good yield and/or with optically high purity.

The present invention is explained by the following examples and comparative examples, but is not limited by those examples.

Example 1

Into a reaction vessel were put potassium phthalimide (50.0g, 0.27mol), benzyltrimethylammonium chloride 5.00g, 0.027mol) and isopropanol (500ml), and the mixture was cooled to 10°C. Thereto was added (R)-epichlorohydrin (74.9g, 0.81mol) and the mixture was stirred for 46 hours while being cooled. The solvent was removed and to the residue was added ethyl acetate (250ml). The mixture was washed with water (250ml), and then ethyl acetate was removed to give crude (R)-glycidylphthalimide (optical purity: 97% e.e.). The crude product was crystallized from ethyl acetate-hexane to give the object (R)-glycidylphthalimide (45.6g, yield: 83%, optical purity: 98%e.e.) as white crystals. mp 100-102C°

Specific rotation $[\alpha]_D^{25} -9.7^\circ$ (c2.0. CHCl₃)

¹HNMR (CDCl₃, 270MHz) δ 2.70 (dd, 1H), 2.81(dd, 1H), 3.21-3.28 (m, 1H), 3.81 (dd, 1H), 3.97 (dd, 1H), 7.27-7.91 (m,

4H)

The optical purity was calculated using HPLC and by its area ratio.

Condition:

Column: CHIRALPAC AD (0.46cm Φ x 25cm L (Daicel Co., Ltd.)

5 Mobile phase: n-hexane/isopropanol (90/10(v/v)

Velocity:1.0ml/min.

Detection: UV220nm

Retention: (S)isomer=17.9min., (R)isomer=25.5min.

10 Example 2

Into a reaction vessel were put potassium phthalimide (5.00g, 27.0mmol), benzyltrimethylammonium chloride (0.50g, 2.70mmol) and tert-butyl alcohol (50ml) and then thereto was added (S)-epichlorohydrin (6.53g, 54.0mmol). The mixture was stirred at 20°C for 24 hours and the solvent was removed. To the residue was added ethyl acetate (30ml), the mixture was washed with water (20ml), and then ethyl acetate was removed to give crude (S)-glycidylphthalimide (3.95g, yield: 72%, optical purity: 97%e.e.) as a white solid.

15

20

Example 3

Into a reaction vessel were put potassium phthalimide (5.00g, 27.0mmol), benzyltrimethylammonium chloride (0.50g, 2.70mmol) and methanol (50ml) and then thereto was added

25

(R)-epichlorohydrin (9.99g, 81.0mmol). The mixture was stirred at 20°C for 15 hours and the solvent was removed. To the residue was added ethyl acetate (30ml), the mixture was washed with water (20ml) and ethyl acetate was removed to give crude (R)-glycidylphthalimide (3.74g, yield: 68%, optical purity: 99%e.e.) as a white solid.

Comparative example 1

Into a reaction vessel were put potassium phthalimide (5.00g, 27.0mmol), benzyltrimethylammonium chloride (0.50g, 2.70mmol) and DMF (50ml) and then thereto was added (R)-epichlorohydrin (5.00g, 54.0mmol). The mixture was stirred at room temperature for 16 hours and the solvent was removed. To the residue was added ethyl acetate (30ml), the mixture was washed with water (20ml) and ethyl acetate was removed to give crude (R)-glycidylphthalimide (4.48g, yield: 82%, optical purity: 63%e.e.) as a white solid.

Comparative examples 2-4 were shown in the following Table 1, which were conducted by changing the solvent according to the method of Comparative example 1.

Table 1

	Solvent	Reaction temp/hr	Yield	Optical purity (%ee)
Comp. ex.2	THF	50°C/44hr	74	87
Comp. ex.3	Toluene	50°C/20hr	n.d	n.d.
Comp. ex.4	1,2-Dichloroethane	30°C/16hr	n.d.	n.d.

Example 4

Into a reaction vessel were put phthalimide (200.0g,
 1.36mol), (S)-epichlorohydrin (226.4g, 2.45mol), anhydrous
 sodium carbonate (14.40g, 0.136mol),
 benzyltrimethylammonium chloride (25.24g, 0.136mol) and
 isopropanol (1.2L) and the mixture was reacted at 25°C for
 22 hours to give crude N-(3-chloro-2-
 hydroxypropyl)phthalimide. The reaction mixture is cooled
 to 15°C and then thereto was dropped a mixture of potassium
 tert-butoxide (183.0g, 1.63mol) and isopropanol (0.8L) over
 a two hours period and the mixture was stirred for 2 hours
 at the same temperature. The solvent was removed and to
 the residue was added ethyl acetate (1.3L). The mixture
 was washed with water (0.65L), and then ethyl acetate was
 removed to give crude (S)-glycidylphthalimide. The crude
 product was crystallized from ethyl acetate-hexane to give
 the object (S)-glycidylphthalimide (206.3g, yield: 75%,
 optical purity: 98%e.e.) as white crystals. mp 99-101°C

Specific rotation $[\alpha]_D^{25} +11.0^\circ$ (c2.0, CHCl_3)

N-(3-Chloro-2-hydroxypropyl)phthalimide

$^1\text{H NMR}$ (CDCl_3 , 270MHz) δ 2.83 (d, 1H), 3.58-3.73 (m, 2H), 3.85-4.01 (m, 2H), 4.13-4.21 (m, 1H), 7.73-7.89 (m, 4H)

5

Example 5

Into a reaction vessel were put phthalimide (2.00g, 13.6mol), (R)-epichlorohydrin (2.52g, 27.2mol), anhydrous sodium carbonate (144mg, 1.36mol), benzyltrimethylammonium chloride (252mg, 1.36mol) and isopropanol (13ml) and the mixture was reacted at room temperature for 15 hours. After removal unreacted epichlorohydrin, to the residue was added isopropanol (13ml) and the mixture is cooled to 20°C . Then thereto was gradually dropped a mixture of potassium tert-butoxide (1.83g, 16.3mol) and isopropanol (7ml) and the mixture was stirred for 1 hour at the same temperature. The solvent was removed and to the residue was added ethyl acetate (13ml). The mixture was washed with water (7ml), and then ethyl acetate was removed to give crude (R)-glycidylphthalimide (2.16g, yield 78%, optical purity 98%e.e.) as white crystals.

In the same manner as Example 5, the following Examples were conducted by using an alkali metal carbonate or an alkali metal hydrogencarbonate with

benzyltrimethylanmonium chloride.

Table 2

	Absolute configuration of epichlorohydrin	Organic solvent	Carbonate or Hydrogencarbonate	Reaction temp. at 1st step	Glycidyl phthalimide	
					Yield	Optical purity (a.c.)
Ex. 6	(S)	t-Butanol	Na_2CO_3	Room temp. /23hr	84%	98%ee (S)
Ex. 7	(R)	Methanol	Na_2CO_3	Room temp. /43hr	61%	98%ee (R)
Ex. 8	(R)	THF	Na_2CO_3	50°C/39hr	81%	97%ee (R)
Ex. 9	(R)	isopropanol	Li_2CO_3	Room temp. /22hr	77%	98%ee (R)
Ex. 10	(R)	isopropanol	K_2CO_3	Room temp. /23hr	75%	98%ee (R)
Ex. 11	(R)	isopropanol	Cs_2CO_3	Room temp. /22hr	70%	98%ee (R)
Ex. 12	(R)	isopropanol	NaHCO_3	Room temp. /25hr	80%	98%ee (R)
Ex. 13	(R)	isopropanol	only Q.A.S.	40°C/22hr	90%	98%ee (R)

Note: in Example 13, only benzyltrimethylanmonium chloride was used as a quaternary ammonium salt (Q.A.S). In Example 7, sodium methoxide (in 28% methanol solution) was used as an alkali metal alkoxide, and in Example 8
5 potassium tert-butoxide, itself was added in several divided portions.

Example 14

Into a reaction vessel were put phthalimide (2.00g,
10 13.6mmol), (R)-epichlorohydrin (2.26g, 24.5mmol), anhydrous sodium carbonate (0.72g, 6.80mmol), tetra n-butylammonium hydrogensulfate (0.46g, 1.36mol) and isopropanol (13ml) and the mixture was reacted at room temperature at 50°C for 23 hours. And then to the residue was gradually added
15 potassium tert-butoxide (1.83g, 16.3mmol) in isopropanol and the mixture is stirred at 20°C for 1 hour. After the reaction, post-treatment was carried out to give crude (R)-glycidylphthalimide (2.21g, yield 80%, optical purity 98%e.e.) as white crystals.

20

Comparative example 5

Into a reaction vessel were put phthalimide (2.00g, 13.6mmol), benzyltrimethylammonium chloride (252mg, 1.36mol) and isopropanol (25mL) and the mixture was ice
25 cooled. Thereto were added potassium tert-butoxide (3.05g,

27.2mmol) in several divided portions and then (R)-epichlorohydrin (2.52g, 27.2mmol). The mixture was reacted at room temperature for 26 hours and then post-treatment was carried out to give crude (R)-glycidylphthalimide (0.45g, yield 17%, optical purity 96%e.e.) as a yellow highly viscous product.

According to the present inventions, there are easily obtainable glycidylphthalimide, especially its optically active compound in good yield with optically high purity and also a N-(3-halo-2-hydroxypropyl)phthalimide, especially its optically active compound in good yield with optically high purity.

The present invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the scope of the present invention, and all such modifications as would be obvious to one skilled in the art are intended to be included within the scope of the following claims.